#### Why have I been given this leaflet?

You have been given this leaflet to provide extra information about some of the tablets you are taking. These medicines are named on the front cover.

These tablets are good for your medical condition. However, if your body becomes short of fluid (dehydration), this medicine can sometimes stop your kidneys from working as they should.

The most common reasons for becoming dehydrated are:

- Vomiting
- Diarrhoea
- High temperatures or fevers
- Not being able to drink normally

If you do develop diarrhoea, vomiting or high fevers, try to compensate by drinking more.

### What should I do with my medicines if I become dehydrated?

#### If you:

- are not able to drink a normal amount of fluid,
- develop diarrhoea or vomiting or
- develop fevers,

you should temporarily stop taking the medicine named on the front of the leaflet. This is to help protect your kidneys.

Please continue to take your other usual medications.

Once you are better and can drink normally, you **should restart** your medicine. For most people this is within 48 hours.

If you remain unwell for longer than this, contact your doctor. It is important to seek medical advice if your symptoms last for more than 48 hours.

## Is there anything else I should do when I am dehydrated?

You <u>can</u> take paracetamol for pain relief or for a high temperature.

**Avoid** anti-inflammatory drugs (a type of pain killer) whilst you're dehydrated. Examples of these medicines are

Ibuprofen, Diclofenac or Naproxen.

To find out more about dehydration and your kidneys, see the NHS Choices website at www.nhs.uk



#### Who is giving you this advice?

This advice comes from kidney specialists and hospital consultants and pharmacists. It aims to try and prevent patients developing kidney problems.

Seek advice from your doctor, pharmacist or nurse If you have any questions about your medicine and its use, or this leaflet.



### What else is important when I am taking this medicine?

If you are coming into hospital for treatment, tell the staff looking after you about this medicine and show them this leaflet. This is particularly important if you are having an operation or some types of scan as your medicines may need to be stopped before your procedure.



25133 V1/Medical Services/TSDFT/03.17/ Review Date 03.19





# Medicines and Your Kidneys

Patient Name	
Medicine(s)	

