

## PATIENT INFORMATION

# Radiotherapy Late Effects Service

Oncology Unit, Torbay Hospital

Radiotherapy late effects are symptoms that can happen any time from 3-6 months to many years after your radiotherapy treatment ended. Or you may find the side effects that you had during treatment never went away.

Many people go on to recover and return to their roles, responsibilities and hobbies after cancer treatment, however, there will be some people who have ongoing issues.

Late effects of radiotherapy can vary between people and depend on which part of the body you had treated. They can include pain or swelling in the area, skin changes, loss of movement or function. If you had your pelvis treated you may have concerns about bowel, bladder, gynaecological or erectile symptoms.

Or, for example, if you had your head and neck treated with radiotherapy you may have concerns about swallowing, dry mouth or thick saliva. Whichever part of

the body you had radiotherapy to, you may have concerns about intimacy and sexual wellbeing.

This is not a complete list of radiotherapy late effects, just some examples.

Radiotherapy late effects can be mild and not cause problems for you, or they can be more severe and affect how you live.

The Radiotherapy Late Effects Service aims to support people to live well with late effects of radiotherapy by listening, offering lifestyle changes and management advice, prescribing medication, providing information or referring to other specialist services.

It is important to realise that it is not always possible to remove radiotherapy late effects altogether.

We hope to reduce the impact the late effects are having and improve your quality of life – this may be by reducing pain, improving bowel or bladder habits, improving function.

This service is here to support and help you with what matters to you in living with these consequences.

You will be offered an appointment with the Late Effects Therapeutic Radiographer and will be given the opportunity to talk about your concerns. A joint action plan to move forward will be agreed between yourself and the radiographer.

You can ask your GP or other healthcare professional involved in your care to refer you to the Late Effects Service. They can do this by using contact details below:

**Contact details:**

Suzanne Hill Late Effects Radiographer, Torbay Oncology Unit, Level 4 Torbay Hospital, Lowes Bridge, Torquay TQ2 7AA

Telephone: 01803 655387

Email: [suzanne.hill@nhs.net](mailto:suzanne.hill@nhs.net)

**Further information**

Macmillan – [Late effects of treatment | Macmillan Cancer Support](#)



---

For further assistance or to receive this information in a different format, please contact the department which created this leaflet.